

Superluminal Noncommutative Photons

Rong-Gen Cai*

Institute of Theoretical Physics, Chinese Academy of Sciences,

P.O. Box 2735, Beijing 100080, China

Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

Abstract

With the help of the Seiberg-Witten map, one can obtain an effective action of a deformed QED from a noncommutative QED. Starting from the deformed QED, we investigate the propagation of photons in the background of electromagnetic field, up to the leading order of the noncommutativity parameter. In our setting (both the electric and magnetic fields are parallel to the coordinate axis x^1 and the nonvanishing component of the noncommutativity parameter is θ^{23}), we find that the electric field has no effect on the propagation of photons, but the velocity of photons can be larger than the speed of light ($c = 1$) when the propagating direction of photons is perpendicular to the direction of background magnetic field, while the light-cone condition does not change when the propagating direction is parallel to the background magnetic field. The causality associated with the superluminal photons is discussed briefly.

*email address: cai@het.phys.sci.osaka-u.ac.jp

Recently much attention has been focused on the noncommutativity of spacetime and its consequences in various aspects. Many interesting phenomena have been found. For example, there exists an intriguing UV/IR mixing [1] in the perturbative dynamics of noncommutative field theories (field theories in a noncommutative spacetime). The degrees of freedom are reduced drastically in the non-planar sector of noncommutative field theories [2–4]. The energy levels of the hydrogen atom and the Lamb shift in the noncommutative quantum electrodynamics (QED) have been found to deviate from those in the usual QED both on the classical and quantum levels, from which a constraint on the noncommutativity parameter of spacetime can be obtained by comparing with experimental data [5]. The noncommutativity of spacetime has also significant effects on the cosmology of the early universe. For instance, it has been shown that the noncommutativity in spatial coordinates can generate magnetic field in the early universe on a horizon scale, and due to its dependence of the number density of massive charged particles, one can trace back the temperature dependence of the noncommutativity scale from the bounds on the primordial magnetic field coming from nucleosynthesis [6]. Further, it was argued that due to the noncommutativity of spacetime, the inflation induced fluctuations become non-Gaussian and anisotropic, and the short distance dispersion relations are modified [7].

By definition, the coordinates in a noncommutative spacetime satisfy

$$[x^\mu, x^\nu] = i\theta^{\mu\nu}, \quad (1)$$

where $\theta^{\mu\nu}$ is an antisymmetric constant matrix and is of dimension of $(length)^2$. The action of a noncommutative field theory can be constructed by replacing the ordinary products between field operators by the $\star$ -products in the action of the corresponding ordinary field theory. The $\star$ -product is defined as follows,

$$f \star g(x) = e^{\frac{1}{2}\theta^{\mu\nu}\partial_{x^\mu}\partial_{y^\nu}} f(x)g(y)|_{y=x}. \quad (2)$$

The Lagrangian of the photon sector of the noncommutative QED is [8]

$$\mathcal{L}_1 = -\frac{1}{4}\hat{F}_{\mu\nu} \star \hat{F}^{\mu\nu}, \quad (3)$$

where $\hat{F}_{\mu\nu} = \partial_\mu \hat{A}_\nu - \partial_\nu \hat{A}_\mu - i[\hat{A}_\mu, \hat{A}_\nu]_\star$. Usually one may start with the action of a certain noncommutative field theory, calculate some physical quantities and then compare with experimental data. However, it is well-known that there is a so-called Seiberg-Witten map [9] between a noncommutative gauge field and an ordinary gauge field. The physical equivalence is guaranteed between the noncommutative gauge field and the ordinary gauge field in the mapping. The Seiberg-Witten map is [9]

$$\hat{A}_\mu = A_\mu - \frac{1}{2}\theta^{\alpha\beta} A_\alpha (\partial_\beta A_\mu + F_{\beta\mu}), \quad (4)$$

to the leading order of $\theta^{\mu\nu}$. Substituting this into (3) and using the $\star$ -product (2), one can obtain [10]

$$\mathcal{L}_2 = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}\theta^{\alpha\beta}F_{\alpha\mu}F_{\beta\nu}F^{\mu\nu} + \frac{1}{8}\theta^{\alpha\beta}F_{\alpha\beta}F_{\mu\nu}F^{\mu\nu}, \quad (5)$$

to the leading order of θ . This Lagrangian can be regarded as the one of the photon sector of the ordinary QED with some correction terms associated with the noncommutativity of spacetime. Physically, the two actions (3) and (5) are equivalent to each other, up to the leading order of θ . Therefore the action (5) plus fermion part could be a good starting point to give some constraints on the realistic noncommutative field theories [11,12].

The form of action (5) is reminiscent of the one-loop effective action of ordinary QED, which is

$$\mathcal{L}_3 = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{\alpha^2}{36m_e^4}(F_{\mu\nu}F^{\mu\nu})^2 + \frac{7\alpha^2}{90m_e^4}F_{\mu\nu}F_{\sigma\tau}F^{\mu\sigma}F^{\nu\tau}, \quad (6)$$

where m_e is the mass of electron, α is the fine structure constant, and last two terms are the Euler-Heisenberg ones. Taking into account the one-loop correction, thirty years ago, Adler [13] showed that the velocity of photons propagating in an anisotropic vacuum given by an external constant uniform magnetic field $\mathbf{B}$, is shifted and depends on the direction of polarizations,

$$v_{\parallel} = 1 - \frac{8}{45}\alpha^2 \frac{\mathbf{B}^2}{m_e^4} \sin^2 \phi, \quad v_{\perp} = 1 - \frac{14}{45}\alpha^2 \frac{\mathbf{B}^2}{m_e^4} \sin^2 \phi, \quad (7)$$

where ϕ is angle between the magnetic field $\mathbf{B}$ and the direction of the propagation. This is just the electromagnetic birefringence phenomenon. Note that here the velocity of photons is always smaller than the speed of light ($c = 1$).

With this in mind, it would be of interest to investigate the propagation of photons within the effective action (5) in the background of electromagnetic field. This is just the aim of this note. Varying the action (5) yields the equation of motion

$$\begin{aligned} \partial_\mu F^{\mu\nu} + \partial_\mu(\theta^{\alpha\beta} F_\alpha{}^\mu F_\beta{}^\nu) - \partial_\mu(\theta^{\alpha\nu} F_{\alpha\lambda} F^{\lambda\mu}) \\ - \frac{1}{2}\partial_\mu(\theta^{\alpha\beta} F_{\alpha\beta} F^{\mu\nu}) + \partial_\mu(\theta^{\alpha\mu} F_{\alpha\lambda} F^{\lambda\nu}) - \frac{1}{4}\partial_\mu(\theta^{\mu\nu} F_{\delta\lambda} F^{\delta\lambda}) = 0. \end{aligned} \quad (8)$$

In order to study the propagation of photons, we expand the field strength using background field method,

$$F^{\mu\nu} = \bar{F}^{\mu\nu} + \tilde{f}^{\mu\nu}, \quad (9)$$

where $\bar{F}_{\mu\nu}$ is the background electromagnetic field. Substituting this into (8) and linearizing the equation in $\tilde{f}^{\mu\nu}$, we obtain the equation describing the propagation of photons

$$\begin{aligned} \partial_\mu \tilde{f}^{\mu\nu} + \theta^{\alpha\beta} \bar{F}_\alpha{}^\mu \partial_\mu \tilde{f}_\beta{}^\nu - \theta^{\alpha\nu} \bar{F}^{\lambda\mu} \partial_\mu \tilde{f}_{\alpha\lambda} - \frac{1}{2}\theta^{\alpha\beta} \bar{F}^{\mu\nu} \partial_\mu \tilde{f}_{\alpha\beta} \\ + \theta^{\alpha\mu} \bar{F}_{\alpha\lambda} \partial_\mu \tilde{f}^{\lambda\nu} + \theta^{\alpha\mu} \bar{F}^{\lambda\nu} \partial_\mu \tilde{f}_{\alpha\lambda} - \frac{1}{2}\theta^{\mu\nu} \bar{F}_{\delta\lambda} \partial_\mu \tilde{f}^{\delta\lambda} = 0, \end{aligned} \quad (10)$$

where the background field has been assumed to be a constant electromagnetic field (namely, $\bar{F}_{\mu\nu}$ is assumed to be a constant). In deriving (10), we dropped some terms like θ times $\partial_\mu \tilde{f}^{\mu\lambda}$ (these are of the order θ^2) since we keep terms up to the leading order of θ only.

In order to investigate the propagation of photons, a simple method is to employ the geometric optic approximation [14], in which one can write

$$\tilde{f}_{\mu\nu} = f_{\mu\nu} e^{i\omega}, \quad (11)$$

where $f_{\mu\nu}$ is a slowly varying amplitude and ω is the rapidly varying phase. The wave vector is $k_\mu = \partial_\mu \omega$. In quantum mechanics, it can be viewed as the momentum of the photons. Thus, from the Bianchi identity, $\partial_\lambda F_{\mu\nu} + \partial_\mu F_{\nu\lambda} + \partial_\nu F_{\lambda\mu} = 0$, one has $k_\lambda f_{\mu\nu} + k_\mu f_{\nu\lambda} + k_\nu f_{\lambda\mu} = 0$. Furthermore, we can write

$$f_{\mu\nu} = k_\mu a_\nu - k_\nu a_\mu, \quad (12)$$

where the vector a_μ can be interpreted as the polarization vector of the photons and satisfies $k_\mu a^\mu = 0$. In this case, the equation (10) reduces to

$$\begin{aligned} & k^2 a^\nu + \theta^{\alpha\beta} \bar{F}_\alpha{}^\mu k_\mu (k_\beta a^\nu - k^\nu a_\beta) - \theta^{\alpha\nu} \bar{F}^{\lambda\mu} k_\mu (k_\alpha a_\lambda - k_\lambda a_\alpha) \\ & - \frac{1}{2} \theta^{\alpha\beta} \bar{F}^{\mu\nu} k_\mu (k_\alpha a_\beta - k_\beta a_\alpha) + \theta^{\alpha\mu} \bar{F}_{\alpha\lambda} k_\mu (k^\lambda a^\nu - k^\nu a^\lambda) \\ & + \theta^{\alpha\mu} \bar{F}^{\lambda\nu} k_\mu (k_\alpha a_\lambda - k_\lambda a_\alpha) - \frac{1}{2} \theta^{\mu\nu} \bar{F}_{\delta\lambda} k_\mu (k^\delta a^\lambda - k^\lambda a^\delta) = 0. \end{aligned} \quad (13)$$

Now it is a position to choose an appropriate background. It has been already shown that a field theory with only nonvanishing space-time component of noncommutative matrix, $\theta^{0i} \neq 0$, is not unitary, while it is unitary if only space-space component $\theta^{ij} \neq 0$ [15]. Furthermore, a field theory with $\theta^{0i} \neq 0$ and $\theta_{\mu\nu} \theta^{\mu\nu} > 0$ is still unitary [16]. In this case, one can change this to the case with only $\theta^{ij} \neq 0$ by an observer Lorentz transformation [16,12]. For simplicity, here we choose

$$\theta^{23} \equiv \theta \neq 0, \quad (14)$$

and other components vanish identically. Further, we suppose that the background electromagnetic field is of the form

$$\bar{F}_{\mu\nu} = -E U_{\mu\nu}^{01} + B U_{\mu\nu}^{23}, \quad (15)$$

where E and B are the strengths of electric and magnetic fields, respectively, and the notation $U_{\mu\nu}^{01}$ has the expression $U_{\mu\nu}^{01} = \delta_\mu^0 \delta_\nu^1 - \delta_\mu^1 \delta_\nu^0$, the other has a similar form. Since one can always choose an appropriate coordinate, in which the electric field and magnetic field are parallel to each other, we therefore assume the background electromagnetic field to be of the form (15). The form (15) implies that both the electric and magnetic fields are parallel to the coordinate axis x^1 . Introducing some linearly independent combinations of momentum components

$$l_\nu = k^\mu U_{\mu\nu}^{01}, \quad m_\nu = k^\mu U_{\mu\nu}^{02}, \quad n_\nu = k^\mu U_{\mu\nu}^{03}, \quad (16)$$

and another dependent combination $p_\nu = k^\mu U_{\mu\nu}^{23}$, and using l_ν , m_ν and n_ν to contract (13), we arrive at

$$\begin{aligned} & k^2(a \cdot v) - 2\theta(a \cdot v)[E(l \cdot p) - B(p \cdot p)] \\ & + \theta(k \cdot v)a^\alpha U_{\alpha\beta}^{23}(El^\beta - Bp^\beta) + \theta[E(l \cdot k) - B(p \cdot k)]a^\alpha v^\beta U_{\alpha\beta}^{23} \\ & - \theta(k \cdot v)p^\alpha a^\lambda (EU_{\alpha\lambda}^{01} - BU_{\alpha\lambda}^{23}) + \theta(k \cdot p)a^\lambda v^\delta (EU_{\lambda\delta}^{01} - BU_{\lambda\delta}^{23}) = 0, \end{aligned} \quad (17)$$

where $v = l, m, n$, respectively. Now we discuss the propagation of photons whose direction is parallel or perpendicular to the one of the background electromagnetic field, respectively.

(a) *The Parallel Case.* Namely, the propagation direction of photons is along x^1 . In this case, we have

$$k^\mu = (k^0, k^1, 0, 0), \quad l^\mu = (k^1, k^0, 0, 0), \quad m^\mu = (0, 0, k^0, 0), \quad n^\mu = (0, 0, 0, k^0), \quad (18)$$

and $p^\mu = 0$. Substituting these into (17) and taking $v = l, m, n$, respectively, it is easy to find that

$$k^2(a \cdot l) = 0, \quad k^2(a \cdot m) = 0, \quad k^2(a \cdot n) = 0, \quad (19)$$

which implies that the light-cone condition of photons does not change not only for the longitudinal polarization ($a \cdot l$), but also for the two transverse polarizations ($a \cdot m$) and ($a \cdot n$). The light-cone condition $k^2 = 0$ indicates that the velocity of photons does not change in this case,

$$v_{x^1} = v_{x^2} = v_{x^3} = 1. \quad (20)$$

(b) *The Perpendicular Case.* Let us suppose that the propagation of photon is along x^3 . In this case, one has

$$k^\mu = (k^0, 0, 0, k^3), \quad l^\mu = (0, k^0, 0, 0), \quad m^\mu = (0, 0, k^0, 0), \quad n^\mu = (k^3, 0, 0, k^0), \quad (21)$$

and $p^\mu = (0, 0, -k^3, 0)$. Substituting these into (17), for the longitudinal polarization ($a \cdot n$) along x^3 , we find

$$k^2(a \cdot n) + 2\theta B(k^3)^2(a \cdot n) = 0, \quad (22)$$

from which we obtain the velocity of photon along x^3

$$v_{x^3} = \left| \frac{k^0}{k^3} \right| \approx 1 + \theta B, \quad (23)$$

to the leading order of θ . For the two transverse polarization $(a \cdot l)$ along x^1 and $(a \cdot m)$ along x^2 , we have

$$(k^2 + 2\theta B(k^3)^2)(a \cdot l) = 0, \quad (k^2 + 2\theta B(k^3)^2)(a \cdot m) = 0. \quad (24)$$

One can see that the light-cone condition is changed, and the velocity of photons becomes

$$v_{x^1} = v_{x^2} \approx 1 + \theta B, \quad (25)$$

up to the leading order of θ .

If the propagation direction of photons is along x^2 , it is easy to show that this case is the same as the one along x^3 . That is, the velocity of photons is also changed in the same way as Eqs. (23) and (25).

From Eqs. (20), (23) and (25), we find some interesting results: in our setup, the background electric field has no effect on the propagation of the noncommutative photon; the light-cone condition does not change for the photons propagating along the direction of background magnetic field; but when propagation direction is perpendicular to the direction of background magnetic field, the velocity of photons is changed and will be larger than the speed of light ($c = 1$) if $\theta B > 0$. (For Dp-branes in the magnetic field, one has $\theta = B^{-1}$ in the decoupling limit [9]. But this cannot be substituted into (23) and (25), since $\theta B \ll 1$ is assumed in our approximation.) Thus, the interesting question arises of whether this superluminal phenomenon is measurable. If we set $\theta \sim (10^4 \text{ GeV})^{-2}$ [5] and $B \sim 1T$, the deviation from the speed of light is then of the order 10^{-24} . It seems beyond the scope that present experiments can reach.

Some additional remarks are in order. First we note that the shift of velocity of photons is a generic phenomenon in nontrivial vacua (for a review see [17]), for example, in

electromagnetic fields [13,18], in gravitational fields [14,18–24], in Casimir-type regions with boundaries [25,26], in finite temperature backgrounds [27] and so on. In the gravitational fields and the Casimir-type regions, the velocity of photons can be larger than the speed of light and it is always smaller than the speed of light in other cases. But our result about the superluminal noncommutative photons is quite different from those, because the shift of velocity of photons in those nontrivial vacua is essentially a *quantum* phenomenon because it appears after taking into account one-loop (and/or) two-loop corrections to the action of usual QED, while our result [see (23) and (25)] is completely a *classical* phenomenon due to the noncommutativity of space, since the action (5) is a classical one without any quantum corrections.

Second, a propagating velocity larger than the speed of light is always associated with a potential violation of causality. For the case of an electromagnetic wave traveling in the vacuum between two parallel conducting plates (Casimir regions), Ben-Menahem [28] showed that the wavefront still travels exactly at the speed of light. So the two-loop effect poses no threat to the causality in QED. In a flat spacetime, in general the establishment of a causal paradox needs at least two condition: spacelike motion and Poincaré invariance. In curved spaces the Poincaré invariance is lost and the principle of equivalence replaces it. In the effective action of QED in curved spaces, however, there are some interaction terms between the electromagnetic field and the spacetime curvature, which violate the principle of equivalence. Therefore the appearance of the superluminal photons in curved spaces is possible and does not necessarily violate the causality. In our case considered in this note, although we are discussing the propagation of photons in a flat spacetime, the Lorentz symmetry is lost due to the noncommutativity of space. As a result, the propagation of superluminal photons is allowed in noncommutative spacetimes and does not threaten the causality.

In summary, we have found a very interesting feature of noncommutative spacetimes: in which the propagation of photons can be faster than the speed of light and it does not necessarily violate the causality.

Note added: The same conclusion has also been obtained independently in [29]. Since this note appeared on the net, I have been informed of many interesting papers, which are related to the conclusion of this note in some sense. In [30,31] it has been shown that solitons in the noncommutative field theories can travel faster than the speed of light along the noncommutative directions. In [32] it has been found that in the noncommutative supersymmetric Yang-Mills the low momenta modes have superluminal group velocity at finite temperature (this is a one-loop effect). However, the propagation of photons in the Born-Infeld theory is always subluminal [33,34]. In addition, the varying speed of light in noncommutative geometry has been used to solve some cosmological problems in the early universe [35].

ACKNOWLEDGMENTS

The author would like to thank Prof. N. Ohta for a reading, Profs. G. Gibbons and R. Jackiw for helpful comments, and especially Prof. R. Jackiw for pointing me out an error in the earlier version. This work was supported in part by a grant from Chinese Academy of Sciences, and in part by the Japan Society for the Promotion of Science and Grants-in-Aid for Scientific Research Nos. 99020 and 12640270.

REFERENCES

- [1] S. Minwalla, M.V. Raamasdonk and N. Seiberg, JHEP **0002**, 020 (2000).
- [2] D. Bigatti and L. Susskind, Phys. Rev. D **62**, 066004 (2000).
- [3] R.G. Cai and N. Ohta, Phys. Rev. D **61**, 124012 (2000).
- [4] W. Fischler, J. Gomis, E. Gorbatov, A. Kashani-Poor, S. Paban and P. Pouliot, JHEP **0005**, 024 (2000).
- [5] M. Chaichian, M.M. Sheikh-Jabbari and A. Tureanu, Phys. Rev. Lett. **86**, 2716 (2001).
- [6] A. Mazumdar and M.M. Sheikh-Jabbari, hep-th/0012363.
- [7] C.S. Chu, B.R. Greene and G. Shiu, hep-th/0011241.
- [8] M. Hayakawa, Phys. Lett. B **478**, 394 (2000).
- [9] N. Seiberg and E. Witten, JHEP **0009**, 032 (1999).
- [10] O.J. Ganor, G. Rajesh and S. Sethi, Phys. Rev. D **62**, 125008 (2000).
- [11] A.A. Bichl, J.M. Grimstrup, L. Popp, M. Schweda and R. Wulkenhaar, hep-th/0102103.
- [12] S.M. Carroll, J.A. Harvey, V.A. Kostelecký, C.D. Lane and T. Okamoto, hep-th/0105082.
- [13] S.L. Adler, Ann. Phys. (New York) **67**, 599 (1971).
- [14] I.T. Drummond and S.J. Hathrell, Phys. Rev. D **22**, 343 (1980).
- [15] J. Gomis and T. Mehen, Nucl. Phys. **B591**, 265 (2000).
- [16] R.G. Cai and N. Ohta, JHEP **0010**, 036 (2000).
- [17] K. Scharnhorst, Annalen Phys. **7**, 700 (1998).
- [18] G.M. Shore, Nucl. Phys. **B460**, 379 (1996).

- [19] R.D. Daniels and G.M. Shore, Nucl. Phys. **B425**, 634 (1994).
- [20] R.D. Daniels and G.M. Shore, Phys. Lett. B **367**, 75 (1996).
- [21] H.T. Cho, Phys. Rev. D **56**, 6416 (1997).
- [22] R.G. Cai, Nucl. Phys. **B524**, 639 (1998).
- [23] A.D. Dolgov and I.D. Novikov, Phys. Lett. B **442**, 82 (1998).
- [24] G.M. Shore, gr-qc/0012063.
- [25] K. Scharnhorst, Phys. Lett. B **236**, 354 (1990).
- [26] G. Barton, Phys. Lett. B **237**, 559 (1990).
- [27] J.I. Latorre, P. Pascual and R. Tarrach, Nucl. Phys. **B437**, 60 (1994).
- [28] S. Ben-Menahem, Phys. Lett. B **250**, 133 (1990).
- [29] Z. Guralnik, R. Jackiw, S.Y. Pi and A. P. Polychronakos, hep-th/0106044.
- [30] D. Bak, K. Lee and J.H. Park, Phys. Rev. **D 63**, 125010 (2001).
- [31] A. Hashimoto and N. Itzhaki, hep-th/0012093.
- [32] K. Landsteiner, E. Lopez and M.H.G. Tytgat, JHEP **0009**, 027 (2000).
- [33] G.W. Gibbons and C.A.R. Herdeiro, Phys. Rev. D **63**, 064006 (2001).
- [34] G.W. Gibbons, hep-th/0104015.
- [35] S.H.S. Alexander and J. Magueijo, hep-th/0104093.